arXiv:hep-ph/0307131v1 9 Jul 2003

# Ground-state baryons in nonperturbative quark dynamics

I.M.Narodetskii and M.A.Trusov
*ITEP, Moscow, Russia*

February 1, 2008

### Abstract

We review the results obtained in an Effective Hamiltonian (EH) approach for the three-quark systems. The EH is derived starting from the Feynman–Schwinger representation for the gauge-invariant Green function of the three quarks propagating in the nonperturbative QCD vacuum and assuming the minimal area law for the asymptotic of the Wilson loop. It furnishes the QCD consistent framework within which to study baryons. The EH has the form of the nonrelativistic three-quark Hamiltonian with the perturbative Coulomb-like and nonperturbative string interactions and the specific mass term. After outlining the approach, methods of calculations of the baryon eigenenergies and some simple applications are explained in details. With only two parameters: the string tension $\sigma = 0.15$ GeV$^2$ and the strong coupling constant $\alpha_s = 0.39$ a unified quantitative description of the ground state light and heavy baryons is achieved. The prediction of masses of the doubly heavy baryons not discovered yet are also given. In particular, a mass of 3660 MeV for the lightest $\Xi_{cc}$ baryon is found by employing the hyperspherical formalism to the three quark confining potential with the string junction.

# 1 Introduction

The heavy flavor physics started with the discovery of the charm quark in 1974. It was followed by the discovery of the beauty quark in 1977 and of the top quark in 1995. Soon after the discovery of the charm quark several charmed baryons and mesons have been identified. The discovery of single charmed or beauty baryons has been followed by the discovery of the $B_c$ meson [1] and recently by the extensive search of doubly charmed baryons [2].

Doubly heavy baryons are baryons that contain two heavy quarks, either $cc$, $bc$, or $bb$. Their existence is a natural consequence of the quark model of hadrons, and it would be surprising if they did not exist. In particular, data from the BaBar and Belle collaborations at the SLAC and KEK B-factories would be good places to look for doubly charmed baryons. Recently the SELEX, the charm hadroproduction experiment at Fermilab, reported a narrow state at $3519 \pm 1$ MeV decaying in $\Lambda_c^+ K^- \pi^+$, consistent with the weak decay of the doubly charmed baryon $\Xi_{cc}^+$ [2]. The candidate is $6.3\sigma$ signal.

The SELEX result was recently critically discussed in [3]. Whether or not the state that SELEX reports turns out to be the first observation of doubly charmed baryons, studying their properties is important for a full understanding of the strong interaction between quarks.

Estimations for the masses and spectra of the baryons containing two or more heavy quarks have been considered by many authors [4]. The purpose of this paper is to present a consistent treatment of the results of the calculation[1] of the masses and wave functions of the doubly heavy baryons obtained in a simple approximation within the nonperturbative QCD. As a by-product, we also report the masses and wave functions for light and heavy baryons.

The paper is organized as follows. In Section 2 we briefly review the Effective Hamiltonian method. In Section 3 we discuss the hyperspherical approach which is a very effective numerical tool to solve this Hamiltonian. In Section 4 our predictions for the ground-state spectra of light and heavy baryons are reported and a detailed comparison of spectra of doubly heavy baryons with the results of other approaches is given. Section 5 contains our conclusions.

# 2 The Effective Hamiltonian in QCD

Starting from the QCD Lagrangian and assuming the minimal area law for the asymptotic of the Wilson loop, the Hamiltonian of the 3q system in the rest frame has been derived. The methodology of the approach has been reviewed recently [6] and so will be sketched here only briefly. The Y-shaped baryon wave function has the form:

$$B_Y(x_1, x_2, x_3, X) = e_{\alpha\beta\gamma} q^{\alpha}(x_1, X) q^{\beta}(x_2, X) q^{\gamma}(x_3, X), \tag{1}$$

where $\alpha$, $\beta$, $\gamma$ are the color indices, $q(x_i, X)$ is the extended operator of the $i^{\text{th}}$ quark at a point $x_i$, and $X = (0, \boldsymbol{X})$ is the equilibrium junction position. The wave function (1) describes the only gauge-invariant configuration possible for baryons. The starting point of the approach is the Feynman–Schwinger representation for the gauge-invariant Green function of the three quarks propagating in the nonperturbative QCD vacuum

---

[1] The preview of this calculation has been done in [5].

$$G(x,y)=\prod_{i=1}^{3}\int_{0}^{\infty} ds_i \int Dz_i \exp(-K_i)\langle \mathcal{W}\rangle_B, \tag{2}$$

where $x = \{x_1, x_2, x_3\}$, $y = \{y_1, y_2, y_3\}$, $z_i = z_i(s_i)$ are the quark trajectories with $z_i(0) = x_i$, $z_i(T) = y_i$, while $s_i$ is the Fock–Schwinger proper time of the $i^{\text{th}}$ quark. Angular brackets mean averaging over background field. The quantities $K_i$ are the kinetic energies of quarks, and all the dependence on the vacuum background field is contained in the generalized Wilson loop $\mathcal{W}$

$$\mathcal{W}=\frac{1}{3!}\varepsilon_{ijk}\varepsilon_{lmn}U_1^{il}U_2^{jm}U_3^{kn}, \tag{3}$$

with

$$U_k = P\exp(ig\int\limits_{C_i} A_\mu(x)dx^\mu), \quad k=1,2,3. \tag{4}$$

Here $P$ denotes the path-ordered product along the path $C_i$ in Fig. 1 where the contours run over the classical trajectories of static quarks. In this figure three quark lines start at junction $X$ at time zero, run in the time direction from 0 to $T$ with the spatial position of quarks fixed and join again in the junction $Y$ at time $T$. There are three planes that are bounded, respectively, by one quark line, two lines connecting junction and quark at $t = 0$ and $t = T$, and the connection line of two junctions. Under the minimal area law assumption, the Wilson loop configuration takes the form

$$\langle \mathcal{W}\rangle_B \propto \exp(-\sigma(S_1+S_2+S_3)), \tag{5}$$

where $S_i$ are the minimal areas inside the contours formed by quarks and the string junction trajectories and $\sigma$ is the QCD string tension.

In Eq. (2) the role of the time parameter along the trajectory of each quark is played by the Fock–Schwinger proper time $s_i$. The proper and real times for each quark are related via a new quantity that eventually plays the role of the dynamical quark mass. The final result is the derivation of the Effective Hamiltonian (EH), see Eq. (6) below.

In contrast to the standard approach of the constituent quark model, the dynamical masses $m_i$ are no longer free parameters. They are expressed in terms of the running masses $m_i^{(0)}(Q^2)$ defined at the appropriate hadronic scale of $Q^2$ from the condition of the minimum of the baryon mass as a function of $m_i$.

Technically, this has been done using the einbein (auxiliary fields) approach, which is proven to be rather accurate in various calculations for relativistic systems. Einbeins are treated as $c$-number variational parameters: the eigenvalues of the EH are minimized with respect to einbeins to obtain the physical spectrum. Such procedure, first suggested in [7], [8], provides the reasonable accuracy for the meson ground states [9].

The method was already applied to study baryon Regge trajectories [7] and very recently for computation of magnetic moments of light baryons [10]. The essential point adopted in [5] and continued in this paper is that it is very reasonable that the same method should also give a unified description for both light and heavy baryons including baryons with two heavy quarks. As in [10] we take as the universal QCD parameter the string tension $\sigma$. We also include the perturbative Coulomb interaction with the frozen strong coupling constant $\alpha_s$.

From experimental point of view, a detailed discussion of the excited $QQ'q$ states is probably premature. Therefore we consider the ground-state baryons without radial and orbital excitations, in which case tensor and spin-orbit forces do not contribute perturbatively. Then only the spin-spin interaction survives in the perturbative approximation. In what follows we disregard the hyper-fine splitting among the baryon masses, then the EH has the following form:

$$H = \sum_{i=1}^{3} \left( \frac{m_i^{(0)^2}}{2m_i} + \frac{m_i}{2} \right) + H_0 + V. \tag{6}$$

Here $H_0$ is the nonrelativistic kinetic energy operator and $V$ is the sum of the perturbative one-gluon-exchange potential $V_C$:

$$V_C = -\frac{2}{3}\alpha_s \cdot \sum_{i<j} \frac{1}{r_{ij}},$$

where $r_{ij}$ are the distances between quarks, and the string potential $V_{\text{string}}$. The baryon mass is given by formula

$$M_B = \min_{m_i} \langle H \rangle + C, \tag{7}$$

where $C$ is the quark self-energy correction calculated in [11]:

$$C = -\frac{2\sigma}{\pi} \sum_i \frac{\eta_i}{m_i} \tag{8}$$

with $\eta = 1$ for $q$-quark[2], $\eta = 0.88$ for $s$-quark, $\eta = 0.234$ for $c$-quark, and $\eta = 0.052$ for $b$-quark.

The string potential calculated in [7] as the static energy of the three heavy quarks was shown to be consistent with that given by a minimum length configuration of the strings meeting in a Y–shaped configuration at a junction $\boldsymbol{X}$:

$$V_{\text{string}}(\boldsymbol{r}_1, \boldsymbol{r}_2, \boldsymbol{r}_3) = \sigma l_{\min}, \tag{9}$$

where $l_{\min}$ is the sum of the three distances between quarks and the string junction point $\boldsymbol{X}$. The Y–shaped configuration was suggested long ago [12], and since then was used repeatedly in many dynamical calculations [13].

# 3 Solving the three-quark equation

## 3.1 Jacobi coordinates

The baryon wave function depends on the three-body Jacobi coordinates

$$\boldsymbol{\rho}_{ij} = \sqrt{\frac{\mu_{ij}}{\mu}} (\boldsymbol{r}_i - \boldsymbol{r}_j), \tag{10}$$

---

[2] Here and throughout the paper $q$ denotes a light quark $u$ or $d$

$$\boldsymbol{\lambda}_{ij} = \sqrt{\frac{\mu_{ij,k}}{\mu}} \left( \frac{m_i \boldsymbol{r}_i + m_j \boldsymbol{r}_j}{m_i + m_j} - \boldsymbol{r}_k \right) \tag{11}$$

$(i, j, k$ cyclic), where $\mu_{ij}$ and $\mu_{ij,k}$ are the appropriate reduced masses

$$\mu_{ij} = \frac{m_i m_j}{m_i + m_j}, \quad \mu_{ij,k} = \frac{(m_i + m_j) m_k}{m_i + m_j + m_k}, \tag{12}$$

and $\mu$ is an arbitrary parameter with the dimension of mass which drops off in the final expressions. The coordinate $\boldsymbol{\rho}_{ij}$ is proportional to the separation of quarks $i$ and $j$ and coordinate $\boldsymbol{\lambda}_{ij}$ is proportional to the separation of quarks $i$ and $j$, and quark $k$. There are three equivalent ways of introducing the Jacobi coordinates, which are related to each other by linear transformations with the coefficients depending on quark masses, with Jacobian equal to unity. In what follows we omit indices $i, j$.

In terms of the Jacobi coordinates the kinetic energy operator $H_0$ is written as

$$H_0 = -\frac{1}{2\mu} \left( \frac{\partial^2}{\partial \boldsymbol{\rho}^2} + \frac{\partial^2}{\partial \boldsymbol{\lambda}^2} \right) = -\frac{1}{2\mu} \left( \frac{\partial^2}{\partial R^2} + \frac{5}{R} \frac{\partial}{\partial R} + \frac{K^2(\Omega)}{R^2} \right), \tag{13}$$

where $R$ is the six-dimensional hyperradius,

$$R^2 = \boldsymbol{\rho}^2 + \boldsymbol{\lambda}^2, \tag{14}$$

$\Omega$ denotes five residuary angular coordinates, and $K^2(\Omega)$ is an angular operator whose eigenfunctions (the hyperspherical harmonics) are

$$K^2(\Omega) Y_{[K]} = -K(K+4) Y_{[K]}, \tag{15}$$

with $K$ being the grand orbital momentum. In terms of $Y_{[K]}$ the wave function $\psi(\boldsymbol{\rho}, \boldsymbol{\lambda})$ can be written in a symbolical shorthand as

$$\psi(\boldsymbol{\rho}, \boldsymbol{\lambda}) = \sum_K \psi_K(R) Y_{[K]}(\Omega).$$

In the hyperradial approximation, which we shall use below, $K = 0$ and $\psi = \psi(R)$. Obviously, this wave function is completely symmetric under quark permutations. Note that the centrifugal potential in the Schrödinger equation for the reduced radial function $\chi(R) = R^{5/2} \psi_K(R)$ with a given $K$

$$\frac{(K+2)^2 - 1/4}{R^2}$$

is not zero even for $K = 0$.

The Coulomb potential is directly expressed in terms of Jacobi coordinates:

$$V_C = -\frac{2\alpha_s}{3} \sum_{i<j} \sqrt{\frac{\mu_{ij}}{\mu}} \cdot \frac{1}{|\boldsymbol{\rho}_{ij}|}, \tag{16}$$

while the corresponding expression for the string potential is more complicated. We will construct it in the next section.

## 3.2 String junction point

Consider the definition of the minimal length string Y–shaped configuration. Let $\varphi_{ijk}$ be the angle between the line from quark $i$ to quark $j$ and that from quark $j$ to quark $k$. One should distinguish two cases. If all $\varphi_{ijk}$ are smaller than 120°, and the equilibrium junction position $\boldsymbol{X}$ coincides with the so-called Torrichelli point of the triangle in which vertices three quarks are situated. If $\varphi_{ijk}$ is equal to or greater than 120°, the lowest energy configuration has the junction at the position of quark $j$.

To find the Torrichelli point, consider a scalar function $\mathcal{L}(\boldsymbol{r})$ of a point inside a triangle △ABC, defined as a sum of distances between this point and the triangle vertices:

$$\mathcal{L}(\boldsymbol{r}) = |\boldsymbol{r} - \boldsymbol{r}_A| + |\boldsymbol{r} - \boldsymbol{r}_B| + |\boldsymbol{r} - \boldsymbol{r}_C|. \tag{17}$$

The position of the minimum of the function $\mathcal{L}$ is calculated from the condition $\dfrac{d\mathcal{L}}{d\boldsymbol{r}} = \boldsymbol{0}$, i.e.:

$$\frac{\boldsymbol{r} - \boldsymbol{r}_A}{|\boldsymbol{r} - \boldsymbol{r}_A|} + \frac{\boldsymbol{r} - \boldsymbol{r}_B}{|\boldsymbol{r} - \boldsymbol{r}_B|} + \frac{\boldsymbol{r} - \boldsymbol{r}_C}{|\boldsymbol{r} - \boldsymbol{r}_C|} = -\boldsymbol{n}_A - \boldsymbol{n}_B - \boldsymbol{n}_C = \boldsymbol{0}, \tag{18}$$

where $\boldsymbol{n}_{A,B,C}$ are the unit vectors from the Torrichelli point directed to the vertices of the triangle. It follows from equation (18) that this condition can be realized only in the case when angles between vectors $\boldsymbol{n}_A$, $\boldsymbol{n}_B$, and $\boldsymbol{n}_C$ are equal to 120°. If $\varphi_{ijk}$ are all smaller than 120°, this point exists and is the unique one. From this point all sides of the triangle are seen at angle of 120°.

The geometrical construction of the Torrichelli point is presented on Fig. 2. One should plot three equilateral triangles △AFB, △BDC, △CEA on the sides of the initial triangle △ABC. It is lightly to prove the following statements:

- The straight lines AD, BE, CF are crossed at a unique point T;
- AD = BE = CF = AT + BT + CT;
- $\angle\mathrm{ATF} = \angle\mathrm{BTF} = \angle\mathrm{BTD} = \angle\mathrm{CTD} = \angle\mathrm{CTE} = \angle\mathrm{ATE} = \dfrac{\pi}{3}$.

Now one can easy obtain an expression for a radius-vector of the Torrichelli point in terms of the lengths $l_i$ of the segments between this point and the $i^{\text{th}}$ quark (segments AT, BT, CT on Fig. 2), and the quark positions $\boldsymbol{r}_i$ (points A, B, C on Fig. 2) [14] :

$$\boldsymbol{X} = \frac{l_2 l_3 \boldsymbol{r}_1 + l_1 l_3 \boldsymbol{r}_2 + l_1 l_2 \boldsymbol{r}_3}{l_2 l_3 + l_1 l_3 + l_1 l_2}. \tag{19}$$

An equivalent expression for $\boldsymbol{X}$ in terms of the center-of-mass position $\boldsymbol{R}_{\text{cm}}$, and vectors $\boldsymbol{\rho}$ and $\boldsymbol{\lambda}$ is [15]

$$\boldsymbol{X} = \boldsymbol{R}_{\text{cm}} + \alpha\boldsymbol{\rho} + \beta\boldsymbol{\lambda}, \tag{20}$$

with

$$\alpha = \frac{1}{2}\sqrt{\frac{\mu}{\mu_{ij}}}\left(\frac{m_j - m_i}{m_i + m_j} - \frac{1}{\sqrt{3}} \cdot \frac{4t + (3 - t^2)\cot\chi}{1 + t^2}\right),$$

$$\beta = \frac{\sqrt{\mu\mu_{ij,k}}}{m_i + m_j} + \sqrt{\frac{\mu}{3\mu_{ij}}} \cdot \frac{\rho}{2\lambda\sin\chi} \cdot \frac{3 - t^2}{1 + t^2},$$

where

$$t = \frac{2\lambda \sin\chi + \sqrt{\frac{3\mu_{ij,k}}{\mu_{ij}}}\rho}{2\lambda\cos\chi + \sqrt{\frac{3\mu_{ij,k}}{\mu_{ij}}} \cdot \frac{m_j - m_i}{m_i + m_j}\rho},$$

and $\chi$ is the angle between $\boldsymbol{\rho}$ and $\boldsymbol{\lambda}$, $0 \le \chi \le \pi$ . It can be easily seen that dependence on $m_i$ in Eq. (20) is apparent and $\boldsymbol{X}$ does not depend on quark masses just as it should be.

Eq. (20) is not very useful for our purposes. What we really need is the explicit expression for $l^2_{\min}$ in terms of the Jacobi coordinates [16]. Introducing the variable $\theta = \arctan(\rho/\lambda)$, $0 \le \theta \le \pi/2$ one obtains for the case $\varphi_{ijk} < 120°$:

$$\begin{aligned} l^2_{\min} = {} & \mu R^2 \cos^2\theta \times \\ & \times \left( \frac{(m_1^3 - m_2^3)\tan^2\theta}{m_1 m_2 (m_1^2 - m_2^2)} + \left( \frac{m_2 - m_1}{m_2 + m_1} \cos\chi + \sqrt{3}\sin\chi \right) \frac{\tan\theta}{m} + \frac{1}{\mu_{12,3}} \right), \end{aligned} \tag{21}$$

where $m^2 = m_1 m_2 m_3 / (m_1 + m_2 + m_3)$. If $m_1 = m_2 = m_3$, this expression coincides with that derived in [7].

If $\varphi_{ijk} > 120°$, $l_{\min} = r_{ij} + r_{jk}$, where

$$\begin{aligned} r_{12} = {} & R\sin\theta \cdot \sqrt{\frac{\mu}{\mu_{12,3}}} \\ r_{13} = {} & R\cos\theta \cdot \sqrt{\frac{\mu}{\mu_{12,3}}} \cdot \sqrt{\frac{m^2}{m_1^2}\tan^2\theta + \frac{2m}{m_1}\tan\theta\cos\chi + 1} \\ r_{23} = {} & R\cos\theta \cdot \sqrt{\frac{\mu}{\mu_{12,3}}} \cdot \sqrt{\frac{m^2}{m_2^2}\tan^2\theta - \frac{2m}{m_2}\tan\theta\cos\chi + 1} \end{aligned} \tag{22}$$

The boundaries corresponding to the condition $\varphi_{ijk} = 120°$ in the $(\chi, \theta)$ plane are:

$$\begin{aligned} \theta_{1(2)}(\chi) &= \arctan(m_{1(2)}(\mp\cos\chi - \sin\chi/\sqrt{3})/m), \\ \theta_3(\chi) &= \arctan(m_2(f(\chi) + \sqrt{f^2(\chi) + 4\varkappa})/2m), \end{aligned} \tag{23}$$

where

$$f(\chi) = (1 - \varkappa)\cos\chi + (1 + \varkappa)\sin\chi/\sqrt{3}, \tag{24}$$

and $\varkappa = m_1/m_2$. These boundaries are shown in Fig. 3 for the case of equal quark masses.

For simplicity, the string junction point is often chosen as coinciding with the center-of-mass coordinate. In this case

$$V_{\text{string}} = \sigma \sum_{(i,j,k)} \frac{1}{m_k} \cdot \sqrt{\mu\mu_{ij,k}} \cdot |\boldsymbol{\lambda}_{ij}| \tag{25}$$

$(i, j, k$ cyclic). Accuracy of this approximation that greatly simplifies the calculations was discussed in [7], [16]. We shall comment on this point later on.

## 3.3 Hyperradial approximation

Introducing the variable $x = \sqrt{\mu} R$ and averaging the interaction $U = V_C + V_{\text{string}}$ over the six-dimensional sphere $\Omega_6$, one obtains the Schrödinger equation for $\chi(x)$:

$$\frac{d^2\chi(x)}{dx^2} + 2\left[E_0 + \frac{a}{x} - bx - \frac{15}{8x^2}\right]\chi(x) = 0. \tag{26}$$

Because the wave function $\psi$ must be finite at the origin $\chi(x) \sim \mathcal{O}(x^{5/2})$ as $x \to 0$. As $x \to \infty$ one can neglect the Coulomb-like and centrifugal terms, and Eq. (26) becomes

$$\frac{d^2\chi(z)}{dz^2} - z\chi(z) = 0, \quad z = (2b)^{1/3}x. \tag{27}$$

This is the familiar Airy equation whose solution $\text{Ai}(z)$ behaves at infinity as

$$\text{Ai}(z) \sim \frac{1}{2}\pi^{-1/2}z^{-1/4}\exp(-\frac{2}{3}z^{3/2}), \quad \text{Re } z \geq\ 0. \tag{28}$$

In Eq. (26) $E_0$ is the ground-state eigenvalue and

$$\begin{aligned} a &= R\sqrt{\mu}\cdot\int V_C(\boldsymbol{r}_1, \boldsymbol{r}_2, \boldsymbol{r}_3)\cdot d\Omega_6 \ = \ \frac{2\alpha_s}{3}\cdot\frac{16}{3\pi}\cdot\left(\sum_{i<j}\sqrt{\mu_{ij}}\right), \\ b &= \frac{1}{R\sqrt{\mu}}\cdot\int V_{\text{string}}(\boldsymbol{r}_1, \boldsymbol{r}_2, \boldsymbol{r}_3)\cdot d\Omega_6. \end{aligned} \tag{29}$$

The expression for the coefficient $b$ can not be obtained analytically except for the equal quark mass system $m_1 = m_2 = m_3 = m$, in which case the straightforward calculation yields:

$$b = \frac{4}{15}\cdot\frac{\sigma}{\sqrt{m}}\cdot\left(-\sqrt{3} + \frac{12\sqrt{2}}{\pi} + \frac{3\sqrt{3}}{\pi}\arccos\frac{1}{5}\right) \approx 1.58\cdot\frac{\sigma}{\sqrt{m}}. \tag{30}$$

On the contrary, in the approximation (25) the coefficient $b$ can be easily found analytically for the case of arbitrary quark masses:

$$b = \sigma\cdot\frac{32}{15\pi}\cdot\left(\sum_{i<j}\frac{\sqrt{\mu_{ij,k}}}{m_k}\right). \tag{31}$$

Let us explain the numerical coefficients in (29), (31) in more details. To this end we introduce the angle $\theta$ as in (21), such that

$$\rho = R\sin\theta, \quad \lambda = R\cos\theta, \quad 0 \leq \theta \leq \frac{\pi}{2}, \tag{32}$$

and write the volume element $d^3\rho d^3\lambda$ as

$$d^3\rho\ d^3\lambda = (4\pi)^2\rho^2\lambda^2 d\rho d\lambda = (4\pi)^2 R^5\sin^2\theta\cos^2\theta dR d\theta. \tag{33}$$

The volume of the six-dimensional sphere is

$$\Omega_6 = (4\pi)^2\int\limits_0^{\frac{\pi}{2}}\sin^2\theta\cos^2\theta d\theta = (4\pi)^2\cdot\frac{\pi}{16} = \pi^3. \tag{34}$$

Then averaging the Coulomb and string terms yields

$$< \frac{1}{\rho} >= \frac{1}{\pi^3} \cdot \frac{1}{R} \cdot (4\pi)^2 \cdot \int\limits_0^{\frac{\pi}{2}} \sin\theta \cos^2\theta \; d\theta = \frac{16}{3\pi} \cdot \frac{1}{R}, \tag{35}$$

$$< \lambda >= \frac{1}{\pi^3} \cdot R \cdot (4\pi)^2 \cdot \int\limits_0^{\frac{\pi}{2}} \sin^2\theta \cos^3\theta \; d\theta = \frac{32}{15\pi} \cdot R. \tag{36}$$

Combining together expressions (16), (25), (29), (35), (36) lead to (29), (31).

## 3.4 Analytic results for light baryons

We can eliminate all dimensional parameters from the equation (26) by a substitution $y = b^{1/3}x$, which leads us to the equation:

$$\frac{d^2\chi}{dy^2} + 2\left(\mathcal{E} - y + \frac{\delta}{y} - \frac{15}{8y^2}\right) = 0 \tag{37}$$

where

$$\mathcal{E} = E_0 b^{-2/3}, \quad \delta = ab^{-1/3}.$$

The eigenvalue of the Eq. (37) can be found using the ordinary perturbation theory, considering the Coulomb term $\left(-\frac{\delta}{y}\right)$ as a small perturbation. This approximation works well for a nucleon containing three light quarks with the running mass equal to zero. In this case there is only one dynamical quark mass $m$. So, the task is greatly simplified and one can obtain analytic expressions for $m$ and $M_B$ via two parameters: $\sigma$ and $\alpha_s$, as expansions in powers of $\alpha_s$.

Omitting the intermediate details outlined in Appendix we quote here the result with accuracy up to $\alpha_s^2$:

$$m = 0.959 \cdot \sqrt{\sigma} \cdot (1 + 0.270\alpha_s + 0.117\alpha_s^2 + \dots), \tag{38}$$

$$M_B = 5.751 \cdot \sqrt{\sigma} \cdot (1 - 0.270\alpha_s - 0.039\alpha_s^2 + \dots) + C, \tag{39}$$

or

$$M_B = 6m \cdot (1 - 0.540\alpha_s - 0.083\alpha_s^2 + \dots) + C, \tag{40}$$

where the constant $C$ has been defined in Eq. (8). As it follows from Eq. (40), the Coulomb-like correction to $M_B$ comprises approximately 20%.

## 3.5 Quasi-classical solution

For the purpose of illustration, the problem is first solved quasi-classically rather than using quantum mechanics. This approach is based on the well known fact that interplay between the centrifugal term and the confining potential produces an effective potential minimum specific for the three-body problem. The numerical solution of (26) for the ground-state eigenenergy may be reproduced on a percent level of accuracy by using the parabolic approximation [17] for the effective potential

$$U(x) = -\frac{a}{x} + bx + \frac{15}{8x^2}.$$

This approximation provides an analytical expression for the eigenenergy. The potential $U(x)$ has the minimum at a point $x = x_0$, which is defined by the condition $U'(x_0) = 0$, i.e.:

$$bx_0^3 + ax_0 - 15/4 = 0\,. \tag{41}$$

Expanding $U(x)$ in the vicinity of the minimum one obtains

$$U(x) \approx U(x_0) + \frac{1}{2}U''(x_0)(x - x_0)^2,$$

i.e., the potential of the harmonic oscillator with the frequency $\omega = \sqrt{U''(x_0)}$. Therefore the ground-state energy eigenvalue is

$$E_0 \approx U(x_0) + \frac{1}{2}\omega\,. \tag{42}$$

## 3.6 Variational solution

Another rather accurate method of solving Eq. (26) is the minimization of the baryon energy using a simple variational Ansätz

$$\chi(x) \sim x^{5/2} e^{-p^2x^2}, \tag{43}$$

where $p$ is the variational parameter. Then using the three-quark Hamiltonian one can get an approximate expression for the ground-state energy: $E_0 \approx \min\limits_p E_0(p)$, where

$$E_0(p) = \langle\chi|H|\chi\rangle = 3p^2 - a \cdot \frac{3}{4} \cdot \sqrt{\frac{\pi}{2}} \cdot p + b \cdot \frac{15}{16} \cdot \sqrt{\frac{\pi}{2}} \cdot \frac{1}{p}. \tag{44}$$

The accuracy of the variational solution will be illustrate in Sect. 4.

## 3.7 Analytical results for $(Qud)$ baryons

For the heavy quarks (Q=c and b) the variation in the values of their masses $m_Q$ is marginal. This is illustrated by the simple analytical results for $(Qud)$ baryons [15]. These results were obtained from the approximate solution of equation

$$\frac{\partial E_0(m_1, m_2, m_3, p)}{\partial p} = 0 \tag{45}$$

where $E_0$ is given by Eq. (44) in the form of expansion in the small parameters

$$\xi = \frac{\sqrt{\sigma}}{m_Q^{(0)}} \quad \text{and} \quad \alpha_s, \tag{46}$$

where $m_Q^{(0)}$ is the heavy-quark running mass.

Omitting the intermediate steps one obtains:

$$E_0 = 3\sqrt{\sigma}\left(\frac{6}{\pi}\right)^{1/4}\left(1 + A\cdot\xi - \frac{5}{3}B\cdot\alpha_s + \dots\right), \tag{47}$$

$$m_q = \sqrt{\sigma}\left(\frac{6}{\pi}\right)^{1/4}\left(1 - A\cdot\xi + B\cdot\alpha_s + \dots\right), \tag{48}$$

$$m_Q = m_Q^{(0)}\left(1 + \mathcal{O}(\xi^2, \alpha_s^2, \alpha_s\xi) + \dots\right), \tag{49}$$

where for the Gaussian variational Ansätz (43)

$$A = \frac{\sqrt{2}-1}{2}\left(\frac{6}{\pi}\right)^{1/4} \approx 0.24,\ \ B = \frac{4+\sqrt{2}}{18}\sqrt{\frac{6}{\pi}} \approx 0.42. \tag{50}$$

Note that the corrections of the first order in $\xi$ and $\alpha_s$ are absent in the expression (49) for $m_Q$. Accuracy of this approximation is illustrated in Table 1 of [15].

# 4 Baryon masses

## 4.1 Quark dynamical masses

We first calculate the dynamical masses $m_i$ retaining only the string potential in the effective Hamiltonian (6). This procedure is in agreement with the strategy adopted in [10]. The masses $m_i$ are then obtained from the equation:

$$\frac{\partial M_B^{(0)}}{\partial m_i} = 0, \tag{51}$$

where

$$M_B^{(0)} = \sum_{i=1}^{3}\left(\frac{{m_i^{(0)}}^2}{2m_i} + \frac{m_i}{2}\right) + E_0(m_1, m_2, m_3; \alpha_s = 0). \tag{52}$$

Then we add the one-gluon-exchange Coulomb potential and solve Eq. (26) to obtain the ground-state eigenvalues $E_0(m_1, m_2, m_3; \alpha_s)$ for a given $\alpha_s$. The physical mass $M_B$ of a baryon is

$$M_B = \sum_{i=1}^{3}\left(\frac{{m_i^{(0)}}^2}{2m_i} + \frac{m_i}{2}\right) + E_0(m_1, m_2, m_3; \alpha_s) + C. \tag{53}$$

We use the values of parameters $\sigma = 0.15$ GeV$^2$ (this value has been found in a recent lattice study [19]) , $\alpha_s = 0.39$, $m_q^{(0)} = 0.009$ GeV ($q = u, d$), $m_s^{(0)} = 0.17$ GeV, $m_c^{(0)} = 1.4$ GeV, and $m_b^{(0)} = 4.8$ GeV. The results for various baryons, obtained using various approximations, are given in Tables 1–3. Table 1 contains the results obtained using the variational solution of Eq. (26) using the approximation (25) for the three-quark potential . In Table 2 the results are shown obtained using the same approximation,

$\boldsymbol{X} = \boldsymbol{R}_{\text{cm}}$, but exact numerical solution of Eq. (26). Table 3 contains the results obtained by the numerical integration of (26) with the genuine three-quark potential in the form (21), (22). Comparing the results of Tables 1 and 2 we observe a good accuracy of the variational solution: the difference between variational and exact results for $M_B$ does not exceed $10-15$ MeV for all baryons from lightest to doubly heavy ones. The approximation $\boldsymbol{X} = \boldsymbol{R}_{\text{cm}}$ leads to a $\sim 5\%$ increase of the coefficient $b$ in (26). As a consequence, this approximation increases the baryon masses by $\sim 70$ MeV (compare the results of Tables 2 and 3.

Note that there is no good theoretical reason why quark masses $m_i$ need to be the same in different baryons. Inspection of Table 1 shows that the masses of the light quarks ($u$, $d$, or $s$) are increased by $\sim 100$ MeV when going from light to heavy baryons. The dynamical masses of light quarks $m_{u,d,s} \sim \sqrt{\sigma} \sim\ 400-500$ MeV qualitatively agree with the results of [20] obtained from the analysis of the heavy-light ground-state mesons.

While studying Table 3 is sufficient to have an appreciation of the accuracy of our predictions, few comments should be added. We expect an accuracy of the baryon predictions to be $\sim 5-10\%$ that is partly due to the approximations employed in the derivations of the Effective Hamiltonian itself [6] and partly due to the error associated with the variational nature of hyperspherical approximation. From this point of view the overall agreement with data is quite satisfactory. E.g. we get $\frac{1}{2}(N+\Delta)_{\text{theory}} = 1144$ MeV vs. $\frac{1}{2}(N+\Delta)_{\text{exp}} = 1085$ MeV ( a 5% increase in $\alpha_s$ would correctly give the $(N,\Delta)$ center of gravity), $\frac{1}{4}(\Lambda+\Sigma+2\Sigma^*) = 1242$ MeV vs. experimental value of 1267 MeV. We also find $\Xi_{\text{theory}} = 1336$ MeV (without hyperfine splitting) vs. $\Xi_{\text{exp}} = 1315$ MeV and $\Xi^c_{\text{theory}} = 2542$ MeV vs. $\Xi^c_{\text{exp}} = 2584$ MeV. On the other hand, our study shows some difficulties in reproducing the $\Omega$-hyperon mass.

## 4.2 Doubly heavy baryons

In Table 4 we compare the spin-averaged masses (computed without the spin-spin term) of the lowest doubly heavy baryons to the predictions of other models [21], [22], [23] as well as variational calculations of [5] for which the center-of-gravity of non-strange baryons and hyperons was essential a free parameter. Most of recent predictions were obtained in a light quark–heavy diquark model [21], [22], in which case the spin-averaged values are $M = \frac{1}{3}(M_{1/2} + 2M_{3/2})$. Note that the wave function calculated in the hyperspherical approximation shows the marginal diquark clustering in the doubly heavy baryons. This is principally kinematic effect related to the fact that in this approximation the difference between the various mean values $\bar{r}_{ij}$ in a baryon is due to the factor $\sqrt{1/\mu_{ij}}$ which varies between $\sqrt{2/m_i}$ for $m_i = m_j$ and $\sqrt{1/m_i}$ for $m_i \ll m_j$. In general, in spite of the completely different physical picture, we find a reasonable agreement within 100 MeV between different predictions for the ground-state masses of the doubly heavy baryons. Our prediction for $M_{ccu}$ is 3.66 GeV with the perturbative hyperfine splitting $\Xi^*_{ccu} - \Xi_{ccu} \sim 40$ MeV. This splitting has been calculated using the Fermi–Breit spin-spin interaction [24]. The change of $\sigma$ to 0.17 GeV$^2$ increases the mass of $\Xi_{cc}$ by $\sim 30$ MeV. Note that the mass of $\Xi_{cc}$ is rather sensitive to the value of the running $c$-quark mass $m_c^{(0)}$, see Fig. 4.

# 5 Conclusions

In this paper, we have outlined a novel approach to baryon spectroscopy which is based on a single framework of the Effective Hamiltonian that is consistent with QCD. This model uses the stringlike picture of confinement and perturbative one-gluon-exchange potential. The main advantage of this work is demonstration of the fact that it is possible to describe all the baryons in terms of the only two parameters inherent to QCD, namely $\sigma$ and $\alpha_s$.

# Acknowledgements

This work was supported in part by NATO grant # PST.CLG.978710 and by RFBR grants ## 03-02-17345, 02-02-17379.

# Appendix

### The approximate calculation of the mass of the three-light-quark system in the hyperspherical formalism

Let us consider the case when each quark has a zero current mass and the constituent mass $m$, which is the same for all three quarks. Then the mass of the system is the minimum of the function $\mathcal{H}(m)$: $\mathcal{M} = \min\limits_m \mathcal{H}(m)$, where[3]

$$\mathcal{H}(m) = \frac{3m}{2} + E\,, \tag{54}$$

and $E$ is the energy level defined from the ordinary Shrödinger equation (see Eq. (26)):

$$\frac{d^2\chi}{dx^2} + 2\left(E + \frac{\beta\sqrt{m}}{x} - \frac{\gamma}{\sqrt{m}}x - \frac{15}{8x^2}\right)\chi = 0\,, \tag{55}$$

where

$$\begin{aligned} \beta &= \frac{16\sqrt{2}}{3\pi}\alpha_s \approx 2.401\cdot\alpha_s, & (56)\\ \gamma &= \frac{32\sqrt{6}}{15\pi}\sigma \approx 1.663\cdot\sigma\,. & (57)\end{aligned}$$

Let us replace a variable $x$ by a dimensionless variable $y$: $y = \left(\frac{\gamma^2}{m}\right)^{1/6} x$. Then:

$$\frac{d^2\chi}{dy^2} + 2\left(\mathcal{E} + \frac{\delta}{y} - y - \frac{15}{8y^2}\right)\chi = 0\,, \tag{58}$$

---

[3] For simplicity we omit here and below the corrections $C$.

where $\mathcal{E}$ and $\delta$ are the dimensionless parameters:

$$\mathcal{E} = E\left(\frac{m}{\gamma^2}\right)^{1/3}, \quad \delta = \beta\left(\frac{m^2}{\gamma}\right)^{1/3}.$$

We will calculate the eigenvalues of the equation (58) using the second-order perturbation theory, considering the Coulomb term $\left(-\frac{\delta}{y}\right)$ in the potential as a small perturbation. The unperturbed equation is:

$$\frac{d^2 f}{dy^2} + 2\left(\lambda - y - \frac{15}{8y^2}\right) f = 0\,. \tag{59}$$

It contains no physical parameters, so its solution is a pure mathematical task. Let us denote the eigenvalues of the equation (59) as $\{\lambda_i\}$:

$$0 < \lambda_0 < \lambda_1 < \dots\,,$$

and the corresponding normalized eigenfunctions as $\{f_i(y)\}$. In what follows we will use the notations:

$$\begin{aligned} \xi &= \int\limits_0^{+\infty} \frac{1}{y} f_0^2(y)dy\,, \quad \xi > 0, \\ \eta &= \sum_{i=1}^{\infty} \frac{\left(\int\limits_0^{+\infty} \frac{1}{y} f_i(y) f_0(y) dy\right)^2}{\lambda_i - \lambda_0}\,, \quad \eta > 0. \end{aligned}$$

The approximate numerical values of these parameters are:

$$\lambda_0 \approx 3.030, \tag{60}$$

$$\xi \approx 0.553, \tag{61}$$

$$\eta \approx 0.028. \tag{62}$$

The ground level of the equation (58) can be approximately calculated as follows:

$$\mathcal{E} \approx \lambda_0 - \delta\xi - \delta^2\eta\,. \tag{63}$$

The small parameter here is the ratio $\frac{\delta\xi}{\lambda_0}$. To estimate this quantity one can solve the task in the zero approximation without the Coulomb term. So,

$$\begin{aligned} \mathcal{E} &\approx \lambda_0, \\ E &\approx \lambda_0\left(\frac{\gamma^2}{m}\right)^{1/3}, \\ \mathcal{H} &\approx \frac{3}{2}m + \lambda_0\left(\frac{\gamma^2}{m}\right)^{1/3}. \end{aligned}$$

$\mathcal{H}$ has a single minimum, defined by an equation $\partial\mathcal{H}/\partial m = 0$, i.e.,

$$\frac{3}{2} - \frac{1}{3}\lambda_0 \gamma^{2/3} m^{-4/3} = 0.$$

Thus $m = \left(\frac{2\lambda_0}{9}\right)^{3/4}\sqrt{\gamma}$, and finally:

$$\frac{\delta\xi}{\lambda_0} = \frac{\beta\xi}{\lambda_0}\left(\frac{m^2}{\gamma}\right)^{1/3} = \frac{2\beta\xi}{3\sqrt{2\lambda_0}} \approx 0.360\alpha_s\,.$$

For $\alpha_s = 0.4$ one has $\frac{\delta\xi}{\lambda_0} \approx 0.14 \ll 1$. This verifies the correctness of using the perturbation theory in this task.

Now, using Eq. (63), we can calculate the constituent mass $m$ and the mass of the state $\mathcal{M}$. For the energy level we have:

$$E = \left(\frac{\gamma^2}{m}\right)^{1/3}\mathcal{E} = \left(\frac{\gamma^2}{m}\right)^{1/3}\lambda_0 - (\gamma m)^{1/3}\beta\xi - m\beta^2\eta\,.$$

It is convenient to use the positive dimensionless parameter $q = m^{1/3}\gamma^{-1/6}$, so that $m = \sqrt{\gamma}q^3$. Then

$$E = \sqrt{\gamma}(\lambda_0 q^{-1} - \beta\xi q - \beta^2\eta q^3)\,,$$

and substituting it to the Eq. (54), we find:

$$\mathcal{H} = \sqrt{\gamma}\left(\lambda_0 q^{-1} - \beta\xi q + \left(\frac{3}{2} - \beta^2\eta\right)q^3\right)\,.$$

$\mathcal{H}$ has a single minimum, defined from the condition: $\left.\frac{\partial\mathcal{H}}{\partial q}\right|_{q=q_0} = 0$, i.e.,

$$\left(\frac{9}{2} - 3\beta^2\eta\right)q_0^2 - \beta\xi - \frac{\lambda_0}{q_0^2} = 0\,. \tag{64}$$

After calculating $q_0$, one can find the constituent mass $m = \sqrt{\gamma}q_0^3$ and the mass of the system $\mathcal{M}$:

$$\mathcal{M} = \mathcal{H}(q_0) = \frac{2\sqrt{\gamma}}{3}\left(\frac{2\lambda_0}{q_0} - \beta\xi q_0\right)\,.$$

The equation (64) can be lightly solved:

$$q_0 = \sqrt{\frac{\beta\xi + \sqrt{\beta^2\xi^2 + 2\lambda_0(9 - 6\beta^2\eta)}}{9 - 6\beta^2\eta}}\,. \tag{65}$$

On expanding the right-hand side of this equation taking $\beta$ as a small parameter

$$q_0 \approx \frac{(2\lambda_0)^{1/4}}{\sqrt{3}}\left(1 + \frac{\xi\beta}{6\sqrt{2\lambda_0}} + \left(\frac{\xi^2}{144\lambda_0} + \frac{\eta}{6}\right)\beta^2\right)\,,$$

we easily get formulas for the constituent mass:

$$m \approx \sqrt{\gamma}\frac{(2\lambda_0)^{3/4}}{3\sqrt{3}}\left(1 + \frac{\xi\beta}{2\sqrt{2\lambda_0}} + \left(\frac{\xi^2}{16\lambda_0} + \frac{\eta}{2}\right)\beta^2\right)\,, \tag{66}$$

and for the state mass:

$$\mathcal{M} \approx \sqrt{\gamma}(2\lambda_0)^{3/4}\frac{2}{\sqrt{3}}\left(1 - \frac{\xi\beta}{2\sqrt{2\lambda_0}} - \left(\frac{\xi^2}{48\lambda_0} + \frac{\eta}{6}\right)\beta^2\right) . \tag{67}$$

Substituting in the equations (66) and (67) numerical values according to formulas (56), (57), (60), (61), (62), we obtain the results quoted in Eqs. (40, (39).

For $\sigma = 0.15$ GeV$^2$, $\alpha_s = 0.4$ one has $m \approx 0.418$ GeV, $\mathcal{M} \approx 1.973$ GeV.

# References

[1] F. Abe *et al.*, Phys. Rev. Lett. **81**, (1998) 2432, hep-ex/9805034.

[2] M. Mattson *et al.*, Phys. Rev. Lett. **89**, 112001 (2002).

[3] V. V. Kiselev and A. K. Likhoded, hep-ph/0208231.

[4] see references [111–124] in hep-ph/0201071.

[5] I. M. Narodetskii and M. A. Trusov, Phys. Atom. Nucl. **65**, 917 (2002) [hep-ph/0104019].

[6] Yu. A. Simonov, Phys. Atom. Nucl. **66**, 338 (2003) [hep-ph/0205334].

[7] M. Fabre de la Ripelle and Yu. A. Simonov, Ann. Phys. (N.Y.) **212**, 235 (1991).

[8] A. Yu. Dubin, A. B. Kaidalov, and Yu. A. Simonov, Phys. Lett. B **323**, 41 (1994).

[9] V. L. Morgunov, A. V. Nefediev, and Yu. A. Simonov, Phys. Lett. B **459**, 653 (1999).

[10] B. O. Kerbikov and Yu. A. Simonov, Phys. Rev. D **62**, 093016 (2000).

[11] Yu.A.Simonov, Phys. Lett. B **515**, 137 (2001).

[12] X. Artru, Nucl. Phys. B **85**, 442 (1975); H. G. Dosch and V. Mueller, Nucl. Phys. B **116**, 470 (1976).

[13] J. Carlson, J. Kogut, and V. R. Pandharipande, Phys. Rev. D **27**, 233 (1983); N. Isgur and J. Paton, Phys. Rev. D **31**, 2910 (1985).

[14] S. Capstick and P. R. Page, Phys. Rev. C **66**, 065204 (2002) [hep-ph/0207027].

[15] I. M. Narodetskii and M. A. Trusov, Nucl. Phys. B (Proc. Suppl.) **115**, 20 (2003) [hep-ph/0209044].

[16] I. M. Narodetskii, A. N. Plekhanov, and A. I. Veselov, JETP Lett. **77**, 58 (2003) [hep-ph/0212235].

[17] Yu. S. Kalashnikova, I. M. Narodetskii, and Yu. A. Simonov, Yad. Fiz. **46**, 1181 (1987).

[18] I. M. Narodetskii and M. A. Trusov, in *Proceedings of the 9th International Conference on the Structure of Baryons (Baryons-2002), 3–8 March, 2002, Newport News, VA, USA* [hep-ph/0304320].

[19] T. T. Takahashi, H. Matsufuru, Y. Nemoto, and H. Suganuma, Phys. Rev. Lett. **86**, 18 (2002).

[20] Yu. S. Kalashnikova and A. Nefediev, Phys. Lett. B **492**, 91 (2000).

[21] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, Phys. Rev. D **62**, 050421 (2000).

[22] D. Ebert, R. N. Faustov, V. D. Galkin, and A. P. Matvienko, Phys. Rev. D **66**, 014008 (2002).

[23] E. Bagan *et al.*, Z. Phys. C **64**, 57 (1994).

[24] A. De Rujula, H. Georgy, and S. L. Glashow, Phys. Rev. D **12**, 147 (1975).

# Tables

**Table 1.** Summary of variational calculations of Eq. (26) for the various baryon states in the approximation $\boldsymbol{X} = \boldsymbol{R}_{\rm cm}$.

| Baryon | $m_1$ | $m_2$ | $m_3$ | $E_0$ | $M_B$ |
|---|---|---|---|---|---|
| $(qqq)$ | 0.372 | 0.372 | 0.372 | 1.433 | 1.221 |
| $(qqs)$ | 0.377 | 0.377 | 0.415 | 1.404 | 1.314 |
| $(qss)$ | 0.381 | 0.420 | 0.420 | 1.377 | 1.405 |
| $(sss)$ | 0.424 | 0.424 | 0.424 | 1.350 | 1.493 |
| $(qqc)$ | 0.424 | 0.424 | 1.464 | 1.178 | 2.538 |
| $(qsc)$ | 0.427 | 0.465 | 1.467 | 1.153 | 2.613 |
| $(ssc)$ | 0.468 | 0.468 | 1.469 | 1.129 | 2.686 |
| $(qqb)$ | 0.446 | 0.446 | 4.819 | 1.093 | 5.909 |
| $(qsb)$ | 0.448 | 0.487 | 4.820 | 1.067 | 5.978 |
| $(ssb)$ | 0.490 | 0.490 | 4.821 | 1.042 | 6.046 |
| $(qcc)$ | 0.459 | 1.498 | 1.498 | 0.914 | 3.712 |
| $(scc)$ | 0.499 | 1.499 | 1.499 | 0.890 | 3.777 |
| $(qcb)$ | 0.477 | 1.524 | 4.834 | 0.793 | 7.021 |
| $(scb)$ | 0.517 | 1.525 | 4.834 | 0.770 | 7.083 |
| $(qbb)$ | 0.495 | 4.854 | 4.854 | 0.606 | 10.260 |
| $(sbb)$ | 0.534 | 4.855 | 4.855 | 0.583 | 10.318 |

**Table 2.** The results obtained from the exact numerical solution of Eq. (26) in the approximation $\boldsymbol{X} = \boldsymbol{R}_{\rm cm}$ for the various baryon states .

| Baryon | $m_1$ | $m_2$ | $m_3$ | $E_0$ | $M_B$ |
|---|---|---|---|---|---|
| $(qqq)$ | 0.372 | 0.372 | 0.372 | 1.427 | 1.212 |
| $(qqs)$ | 0.376 | 0.376 | 0.415 | 1.398 | 1.306 |
| $(qss)$ | 0.381 | 0.419 | 0.419 | 1.370 | 1.397 |
| $(sss)$ | 0.423 | 0.423 | 0.423 | 1.344 | 1.485 |
| $(qqc)$ | 0.424 | 0.424 | 1.464 | 1.170 | 2.530 |
| $(qsc)$ | 0.426 | 0.465 | 1.466 | 1.146 | 2.604 |
| $(ssc)$ | 0.467 | 0.467 | 1.468 | 1.122 | 2.677 |
| $(qqb)$ | 0.445 | 0.445 | 4.820 | 1.085 | 5.900 |
| $(qsb)$ | 0.448 | 0.487 | 4.820 | 1.059 | 5.970 |
| $(ssb)$ | 0.488 | 0.488 | 4.820 | 1.035 | 6.037 |
| $(qcc)$ | 0.458 | 1.497 | 1.497 | 0.905 | 3.702 |
| $(scc)$ | 0.497 | 1.498 | 1.498 | 0.882 | 3.767 |
| $(qcb)$ | 0.475 | 1.523 | 4.833 | 0.784 | 7.010 |
| $(scb)$ | 0.515 | 1.523 | 4.837 | 0.760 | 7.072 |
| $(qbb)$ | 0.490 | 4.850 | 4.850 | 0.596 | 10.245 |
| $(sbb)$ | 0.530 | 4.856 | 4.856 | 0.571 | 10.303 |

**Table 3.** The same as in Table 2 but for an exact treatment of the string-junction point.

| Baryon | $m_1$ | $m_2$ | $m_3$ | $E_0$ | $M_B$ |
|---|---|---|---|---|---|
| $(qqq)$ | 0.362 | 0.362 | 0.362 | 1.392 | 1.144 |
| $(qqs)$ | 0.367 | 0.367 | 0.407 | 1.362 | 1.242 |
| $(qss)$ | 0.371 | 0.411 | 0.411 | 1.335 | 1.336 |
| $(sss)$ | 0.415 | 0.415 | 0.415 | 1.307 | 1.426 |
| $(qqc)$ | 0.406 | 0.406 | 1.470 | 1.142 | 2.464 |
| $(qsc)$ | 0.409 | 0.448 | 1.471 | 1.116 | 2.542 |
| $(ssc)$ | 0.452 | 0.452 | 1.473 | 1.090 | 2.621 |
| $(qqb)$ | 0.425 | 0.425 | 4.825 | 1.054 | 5.823 |
| $(qsb)$ | 0.429 | 0.469 | 4.826 | 1.026 | 5.903 |
| $(ssb)$ | 0.471 | 0.471 | 4.826 | 1.000 | 5.975 |
| $(qcc)$ | 0.444 | 1.494 | 1.494 | 0.876 | 3.659 |
| $(scc)$ | 0.485 | 1.496 | 1.496 | 0.851 | 3.726 |
| $(qcb)$ | 0.465 | 1.512 | 4.836 | 0.753 | 6.969 |
| $(scb)$ | 0.505 | 1.514 | 4.837 | 0.729 | 7.032 |
| $(qbb)$ | 0.488 | 4.847 | 4.847 | 0.567 | 10.214 |
| $(sbb)$ | 0.526 | 4.851 | 4.851 | 0.544 | 10.273 |

**Table 4**. Comparison of various predictions for ground-state masses (in units of GeV) of doubly heavy baryons.

| Baryon | Ref. [16] | Ref. [5] | Ref. [21] | Ref. [22] | Ref. [23] |
|---|---|---|---|---|---|
| $\Xi_{cc}$ | 3.66 | 3.69 | 3.57 | 3.69 | 3.70 |
| $\Omega_{cc}$ | 3.73 | 3.86 | 3.66 | 3.84 | 3.80 |
| $\Xi_{cb}$ | 6.97 | 6.96 | 6.87 | 6.96 | 6.99 |
| $\Omega_{cb}$ | 7.03 | 7.13 | 6.96 | 7.15 | 7.07 |
| $\Xi_{bb}$ | 10.21 | 10.16 | 10.12 | 10.23 | 10.24 |
| $\Omega_{bb}$ | 10.27 | 10.34 | 10.19 | 10.38 | 10.34 |

# Figure captions

**Figure 1.** Three-lobes Wilson loop.

**Figure 2.** The geometrical construction of the Torrichelli point T for an arbitrary triangle $\triangle$ABC. The triangles $\triangle$AFB, $\triangle$BDC, $\triangle$CEA are equilateral.

**Figure 3.** The four regions in the $(\chi, \theta)$ plane corresponding to $\varphi_{ijk} \geq 120^\circ$ and $\varphi_{ijk} \leq 120^\circ$ for the case of equal quark masses.

**Figure 4.** Mass of $\Xi_{cc}$ as a function of the running $c$-quark mass for $\sigma = 0.15$ GeV$^2$ and $\sigma = 0.17$ GeV$^2$. The masses are given in GeV. Bold points refer to the case $m_c^{(0)} = 1.4$ GeV.



**Fig.1.**



**Fig.2.**


π/2
cosθ$_{231}$ < -1/2
θ$_3$
θ$_2$
θ$_1$
cosθ$_{ijk}$ > -1/2
cosθ$_{123}$ < -1/2
cosθ$_{312}$ < -1/2
0
π/3
2π/3
π

**Fig.3.**


$M_{\Xi_{cc}}$, GeV
3.4
3.5
3.6
3.7
3.8
1.3
1.35
1.4
1.45
$m_c^{(0)}$, GeV
$\sigma = 0.17$ GeV$^2$
$\sigma = 0.15$ GeV$^2$

**Fig.4.**